



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

both by Principal Forbes and Sir William Thomson, the author applies himself solely to trace the existence of other cycles than the ordinary annual one, in the rise and fall of the different thermometers.

Of such cycles, and of more than one year's duration, he considers that he has discovered three; and of these the most marked has a period of 11·1 years, or practically the same as Schwabe's numbers for new groups of solar spots. Several numerical circumstances, however, which the author details, show that the sun-spots cannot be the actual cause of the observed waves of terrestrial temperature, and he suggests what may be, concluding with two examples of the practical use to which a knowledge of the temperature cycles as observed may at once be turned, no matter to what cosmical origin their existence may be owing.

II. "On the Constituent Minerals of the Granites of Scotland, as compared with those of Donegal." By the Rev. SAMUEL HAUGHTON, F.R.S., M.D. Dubl., D.C.L. Oxon., Fellow of Trinity College, Dublin. Received March 31, 1870.

During the past summer (1869) I completed my investigation of the constituent minerals of the Scotch Granites, and secured specimens, from the analysis of which I obtained the following results:—

#### I. *Orthoclase*.

	No. 1.	No. 2.	No. 3.	No. 4.
Silica . . . . .	65·40	64·44	64·48	64·48
Alumina . . . . .	19·04	18·64	20·00	20·00
Peroxide of iron . . .	trace.	0·80	none.	none.
Lime . . . . .	0·22	0·66	1·01	0·78
Magnesia . . . . .	trace.	trace.	trace.	none.
Soda . . . . .	3·63	2·73	1·72	2·19
Potash . . . . .	11·26	12·15	12·81	12·10
Water . . . . .	0·20	0·80	0·64	0·08
	<hr/> 99·75	<hr/> 100·22	<hr/> 100·66	<hr/> 99·63

No. 1. Stirling Hill, Peterhead. Occurs in an eruptive Granite, in veins, in well-developed reddish-pink opaque crystals, encrusted with crystals of Albite.

No. 2. Rubislaw, Aberdeen. Large beautiful reddish-pink opaque crystals in veins, associated with white Mica. The Granite of Rubislaw is of metamorphic origin, and different in character from the eruptive Granite of Peterhead. No Albite has been found in it.

No. 3. Peterculter, Aberdeen. In metamorphic Granite; white, translucent, large crystals.

No. 4. Callernish, extreme west of Lewis. In metamorphic Granite; in large grey crystals, with a slight shade of pink, translucent.

The oxygen ratio of these felspars is as follows:—

	No. 1.	No. 2.	No. 3.	No. 4.
Silica .....	33·956	33·456	33·478	33·477
Alumina &c. ..	8·898	8·950	9·348	9·348
Lime .....	0·061	0·187	0·286	0·221
Soda .....	0·929	0·699	0·440	0·561
Potash .....	1·908	2·059	2·171	2·051
	<hr/> 45·752	<hr/> 45·351	<hr/> 45·723	<hr/> 45·658

From this Table we find the oxygen ratios:—

	No. 1.	No. 2.	No. 3.	No. 4.
Silica .....	11·37	11·35	11·55	11·82
Peroxides .....	3·06	3·04	3·22	3·30
Protoxides ....	1·00	1·00	1·00	1·00

The Granites of central and western Scotland are metamorphic rocks, like those of Donegal and Norway, with which they are geologically identical; and truly eruptive Granite occurs at only a few localities, as, for example, near Peterhead.

The second felspar associated with Orthoclase in the Metamorphic Granites is Oligoclase, as in Donegal; while the second felspar associated with Orthoclase in the eruptive Granites is Albite, as in Mourne, Leinster, and Cornwall. The fact thus indicated by the Scotch Granites is completely in accordance with the mode of occurrence of Oligoclase and Albite in the Irish Granites.

## II. *Oligoclase.*

	No. 1.	No. 2.
Silica .....	62·00	61·88
Alumina .....	23·20	24·80
Magnesia .....	—	Trace.
Lime .....	4·71	4·93
Soda .....	9·20	8·12
Potash .....	0·43	0·98
	<hr/> 99·54	<hr/> 100·71

No. 1. This Oligoclase occurs in the Granite of Craigie-Buckler, near Aberdeen; it is white and opaque, and so much resembles Cleavelandite in appearance as to have been mistaken for that variety of Albite; its analysis proves it to be Oligoclase. The crystals do not exhibit striation.

No. 2. From the Granite of Rhiconich, in the west of Sutherlandshire;

it is greyish white, semitranslucent, in large striated crystals, and resembles the Oligoclase of Ytterby, in Sweden.

The oxygen ratios of the Oligoclase are as follow :—

	No. 1.	No. 2.
Silica .....	32·191	32·128
Alumina .....	10·843	11·590
Lime .....	1·339	1·400
Soda .....	2·360	2·082
Potash .....	0·072	0·165
	<hr/> 46·805	<hr/> 47·365

Hence we obtain :—

	No. 1.	No. 2.
Silica .....	8·54	8·82
Peroxides .....	2·88	3·18
Protoxides .....	1·00	1·00

These oxygen ratios prove the felspars to be Oligoclase.

### III. *Albite*.

Silica .....	68·00
Alumina .....	20·00
Lime .....	0·35
Magnesia .....	trace.
Soda .....	10·88
Potash .....	0·68
	<hr/> 99·91

This Albite occurs at Stirling Hill, near Peterhead, in eruptive Granite, and is found associated with red Orthoclase in veins; it encrusts the large crystals of Orthoclase, and is semitranslucent, and is generally stained on the surface by peroxide of iron.

### *Oxygen Ratios.*

Silica .....	35·306	.....	11·77
Alumina .....	9·348	.....	3·11
Lime .....	0·099	}	1·00
Soda .....	2·790		
Potash .....	0·114		

This mineral is evidently a typical Albite.

There are two kinds of Mica found in the Scotch Granites, and both Micæ resemble very closely the corresponding minerals of the Donegal Granites.

IV. *White Mica.*

Silica .....	44·40
Fluosilicon ( $\text{Si F}_3$ ).....	0·16
Alumina .....	37·36
Peroxide of iron .....	2·04
Lime .....	0·78
Magnesia .....	0·57
Soda .....	0·93
Potash .....	9·87
Protoxide of manganese .....	0·24
Water .....	1·84
	<hr/>
	98·19

The specimen of Mica here analyzed came from veins in the Granite quarry of Rubislaw, near Aberdeen, and occurs in large plates, associated with red Orthoclase. It was carefully examined for lithia, but no trace of this alkali could be found in it.

The angles of the rhombic plates were  $60^\circ$  and  $120^\circ$  exactly, and the angle between its optic axes was found to be  $72^\circ 30'$ .

The black Mica in large crystals is very rare, but it seems abundantly disseminated in minute scales through most of the Scotch Granites. The following analysis was made on specimens found near Aberdeen by Prof. Nicol, and kindly forwarded to me by him, for the purposes of this paper:—

V. *Black Mica.*

Silica .....	36·50
Alumina .....	16·50
Peroxide of iron .....	18·49
Lime .....	1·11
Magnesia .....	7·44
Soda .....	0·92
Potash .....	8·77
Protoxide of iron .....	6·76
Protoxide of manganese .....	1·80
Water .....	1·60
	<hr/>
	99·89

This mica was carefully examined for fluorine, and found not to contain any.